Journal of Scientific Research and Reports



Volume 30, Issue 11, Page 483-493, 2024; Article no.JSRR.126176 ISSN: 2320-0227

# Influence of Potting Mixture and Sowing Orientation on Physiological Parameters of *Balanites roxburghii* (Linn.)

# Basavaraj S Ganigara <sup>a++\*</sup>, Doddabasawa <sup>b#</sup>, Hanumanthappa Dasanahalli <sup>c++</sup>, Shakuntala N.M <sup>d†</sup>, Umesh Hiremath <sup>e‡</sup> and Maruti.K <sup>e‡</sup>

<sup>a</sup> Department of Seed Science & Technology, College of Agriculture, Bheemarayanagudi, University of Agricultural Sciences, Raichur, Karnataka (585287), India.
 <sup>b</sup> Department of Agro Forestry and Environment, College of Agriculture, Bheemarayanagudi, University of Agricultural Sciences, Raichur, Karnataka, (585287), India.
 <sup>c</sup> Agricultural Research Station, Gudadalli, UAS, Raichur, India.
 <sup>d</sup> Department of Seed Science & Technology, College of Agriculture, Raichur, UAS, Raichur, India.
 <sup>e</sup> Seed Unit, National Seed Project, UAS, Raichur, India.

# Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

# Article Information

DOI: https://doi.org/10.9734/jsrr/2024/v30i112576

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/126176

> Received: 01/09/2024 Accepted: 02/11/2024 Published: 08/11/2024

**Original Research Article** 

*Cite as:* Ganigara, Basavaraj S, Doddabasawa, Hanumanthappa Dasanahalli, Shakuntala N.M, Umesh Hiremath, and Maruti.K. 2024. "Influence of Potting Mixture and Sowing Orientation on Physiological Parameters of Balanites Roxburghii (Linn.)". Journal of Scientific Research and Reports 30 (11):483-93. https://doi.org/10.9734/jsrr/2024/v30i112576.

<sup>++</sup> Assistant Professor of SST;

<sup>#</sup> Associate Professor of Forestry;

<sup>&</sup>lt;sup>†</sup> Professor of SST;

<sup>&</sup>lt;sup>‡</sup> Assistant Professor of Seed Science & Technology;

<sup>\*</sup>Corresponding author: E-mail: bganiger07@gmail.com;

# ABSTRACT

Balanites roxburghii (Linn.) species has been classified under the category of endangered plant species due to excessive exploitation for a variety of purposes, which has necessitated conservation of this tree species, and hence a study was conducted to investigate the effect of potting mixture and sowing orientation on early growth performance and vigour. Seeds were subjected to seven potting mixtures: M1: Soil:Sand:FYM 2:1:1, M2: Soil:Cocopeat:FYM 2:1:1,M3: Soil:Cocopeat:Vermicompost 2:1:1, M4: Soil:Paddy Husk:FYM: 2:1:1, M5: Soil:Paddy Husk:Vermicompost: 2:1:1, M6: Soil: Saw Dust:FYM 2:1:1, and M7: Soil:sawdust: Vermicompost 2:1:1. The seeds were sown in three orientations. S1: Seeds laid vertically with stalk end downwards; S2: Seeds laid horizontally and S3: Seeds laid vertically with stalk end upwards. Physiological parameters considered for assessment were seed germination, days taken for germination, seedling length, and seedling vigor index. A completely randomized design was used for the experiment. The results showed that there is a significant difference among seven different potting mixtures and three orientations. At the end of 8 weeks, the seed is sown horizontally in the potting mixture Soil:cocopeat:vermicompost (2:1:1) has recorded the germination percentage of 71.67%, 37 days taken for seed germination, highest seedling length (64 cm), and highest seedling vigour index (4561). The high-vigor seedlings may survive even in unfavorable conditions, with high survival rate compared to others. The study recommended that to obtain high-vigor Balanites roxburghii seedlings, seeds may be sown in potting mixture soil: cocopeat: vermicompost (2:1:1) with horizontal orientation.

Keywords: Potting mixture; orientation; seedling vigour index; seed germination.

# 1. INTRODUCTION

Balanites roxburghii (Linn.) is commonly called as 'Desert Date' in English belongs to family of Zygophyllaceae. It is one of the most neglected and underutilized plant species found in dry areas of Africa and South Asia (Daya & Vaghasiya, 2011). In India, it is naturally found in Rajasthan, Gujarat, Madhya Pradesh and Deccan Plateau (Chhanval et al., 2023). This plant grows in harsh climate (arid and semi-arid) and wide variety of soil ranging from sandy loamy soil to deep black soil. The plant grows as a thorny shrub to small tree and is being locally used for multiple utility especially for its medicinal value. The fruits, pulp, seed, leaves, bark and roots have antimicrobial, anticancer, antiviral, antidiabetic, antioxidant, anti-inflammatory and other properties used locally to cure many diseases such as jaundice, intestinal worm infection, wounds, malaria, syphilis, epilepsy, dysentery, constipation, diarrhea, hemorrhoid, stomach aches, asthma, and fever (Al-Thobaiti and Zeid, 2018; Saboo et al. 2014; Hammiche and Maiza, 2006). The leaves and fruits are used as a fodder for livestock (Orwa et al., 2009). The diosgenin content in roots and fruits are used to make oral contraceptive, sex hormone and steroids (Chapagain et. al., 2009). Young shoots and leaves are used vegetables by the tribes and In Africa ripen fruits are eaten raw or after sun dried. Further, in India pulp of the fruit is

used as detergent to wash the cloths and wood is used to make agricultural implements and as a fuel wood and it is also being assisted on the boundary of agriculture land as live fence since as it is a thorny species with small crown and height.

This species is one of the natural compositions of dry evergreen or thorn forest of semi-arid climate. However, in recent past the population of this species is dwindling due to various factors such as poor germination, biotic pressure, forest fire and reduced ethnobotanical usage.

The species is mainly regenerated through seeds; however, the seeds of this species are of varying in size, weight and germination percentage. As fruits are of brittle and smooth mesocarp and consists of oil and fibers endocarp. Hence, freshly sown seeds have higher germination compared to too late sowing and it studies indicated the species do not require seed treatment. However, treating seeds with cold and hot water reported enhanced regeneration whereas, minimal work is done on the potting mixture which is most important for raising seedlings in nursery for afforestation and conservation and similarly on the plant orientation of sowing seeds. Keeping this in view the present investigation was undertaken to know the influence of potting mixture and orientation of sowing seed on physiological traits.

### 2. MATERIAL AND METHODS

The present investigation on influence of potting mixture and orientation of seeds on physiological traits of Balanites roxburghii were carried out in and around Raichur district of Karnataka, India, which falls under North Eastern Dry Zone (Zone-II) of Karnataka Agroclimatic Zones and study site is located between 16°.15 N latitude and 77°.20 E longitude and at 398.37 m above mean sea level. Average monthly temperature of this area normally around 25°C (January) to 35°C (July) with a mean annual rainfall of 680 mm and climate of the study area is of semi-arid with mild temperature in the winter and high harsh temperature in summer. The soils are mainly medium to deep black soils in major part and also red loamy and sandy loamy in few pockets.

The ripen fruits were collected in the month of January from well grown trees from nearby area and on the boundary of agricultural lands and fruits brought to the laboratory and were dried under shades for 15 days later fruits soaked in the cold water for 24 hours and were de-pulped and were sown in the pots (Ahmed et al., 2017). The experiment was carried out with CRD in pots with two factors of which 7 main factors with different potting mixture and 3 sub factors of seed orientation. Further, 25 de-pulped seeds were sown for each factor and were replicated into four times. The details of the main and sub factors are given below.

#### Factor I: Potting media

- M1: Soil: Sand: FYM 2:1:1
- M2: Soil: Coco peat: FYM 2:1: 1
- M3: Soil: Coco peat: Vermicompost 2:1:1
- M4: Soil: Paddy husk : FYM : 2:1:1
- M5: Soil : Paddy husk : Vermicompost : 2:1:1
- M6: Soil : Saw dust : FYM 2:1:1
- M7: Soil : saw dust : vermicompost 2:1:1

#### **Factor II: Orientation**

- S1: Seed laid vertically with stalk end downwards
- S2: Seeds laid horizontally
- S3: Seeds laid vertically with stalk end upwards

The experiment was continuously monitored and regular watering was done to keep the adequate moisture for 8 weeks. The data on germination was observed daily and seeds were considered germinated on emergence of healthy plumule through media and data obtained were subjected to assess germination, seedling length and seedling vigor index using the following equations. Then, data were subjected to statistical analysis using MS-Excel at P level of 0.01.

Table 1.	Nitrogen,	phosphorus	s and p	otash
	content in	n different m	edia	

Media	N %	P%	K%
Cocopeat	0.41	0.81	1.30
FYM	0.50	0.30	0.50
Rice husk	0.31	0.19	0.22
Vermicompost	1.25	1.14	1.19

**Seed germination assessment:** The data on germination was recorded on weekly basis for a period of eight weeks. The germination percentage was calculated by the following formula:

Seed Germination (%) =  $\frac{\text{Germinated seeds}}{\text{Total number of seeds}} \times 100$ 

**Seedling length (cm):** Ten normal seedlings were selected at random from each treatment. The seedling length was measured from the shoot tip to root tip of the seedling and the mean was expressed in centimeters (cm).

**Seedling vigour index-I:** The seedling vigour index-I was calculated as per the formula given by Abdul-Baki and Anderson (1973) expressed as whole number.

Seedling vigour index-I = Germination (%) x Mean seedling length (cm)

#### 3. RESULTS AND DISCUSSION

#### 3.1 Seed Germination

The investigation of influence of potting mixture and orientation of seeds on physiological traits of Balanites revealed significant difference with pottina mixture and seed orientation. Significantly, higher seed germination was noticed in M1 potting mixture Soil : Sand : FYM (2:1:1) (69.00%), followed by Soil: Coco peat: Vermicompost M3 (2:1:1) (59.56%). Whereas significantly lower germination was recorded in M4 Soil: Paddy husk : FYM (2:1:1) (41.89%) (Table 2). This could be attributed that standard ratio followed for nursery raising seedlings as sand helps for better aeration followed by adequate nutrients from soil and FYM helped the germination as the seed of Balanites has more surface area and bigger in size and lower germination in treatments could be due to lack of aeration. Similarly, Akanbi et al. (2002) reported that, for better germination of seeds needs adequate aeration and nutrients.

Among the different seed sowing orientations, significantly higher seed germination was observed with seed oriented vertically with stalk end upwards (67.76%) and seed laid horizontally (61.14%) respectively. However, significantly, lower seed germination was recorded in seed oriented vertically with stalk end downwards (31.19%). Since orienting seeds with stalk end upwards and laying seeds horizontally results in less difference in seed germination percentage. It is better to sow the seeds horizontally since this is the normal orientation of seeds/fruits, when fall on the ground. Also, orienting seeds with stalk end with upwards by inexperienced persons can lead to sowing the seeds in stalk end upwards. Seeds must be placed in a position that allowed the uptake of water and other environmental variables required for germination (Bowers & Hayden, 1972). These results are line with the findings of Elffel (2012), El-Nour & Kalislo (1995) and Wakawa & Suleiman (2022) in Balanities aegyptiaca.

The study on interaction between potting mixture and orientation revealed significant difference. Significantly, higher seed germination (84.33) was recorded in Soil:Sand:FYM (2:1:1) with Seeds laid vertically with stalk end upwards followed by the potting mixture Soil : Sand : FYM (2:1:1) with seeds laid horizontally (81.00%). Whereas, significantly, lower seed germination percentage was recorded inSoil: Paddy husk : FYM (2:1:1) with Seed laid vertically with stalk end downwards (23.33%). This might be due to better aeration in potting mixture with sand and adequate nutrients as well as orientation of the seed. If seeds sown vertically with stalk downwards may die due to lack of good aeration and needs lot of energy to produce the plumules which comes from the stalk. Hence, to get higher germination it is better to sow the seeds vertically keeping stalk at upwards.

# 3.2 Days to Germination

The study on time taken to complete germination noticed significant difference with potting

mixture and orientation (Table 3). Significantly, minimum numbers of days taken to complete germination were observed in soil:Sand:FYM (28.78 days) followed by Soil:Cocopeat :FYM (34.44 days) and Soil:Cocopeat: Vermicompost (37.44 days). However, significantly maximum number of days were taken complete the germination was recorded in the potting mixture Soil:Paddy husk: Vermicompost (45.67 days). This may be attributed better aeration and adequate nutrients in potting mixture with sand as compared to without sand.

Similarly, significant difference in time taken to complete germination was noticed with sowing orientation. Significantly, minimum number of days taken to complete germination were recorded in seeds sown vertically with stalk upward germinate (31.33 days) followed by horizontal sowing. However, significantly, maximum number of days taken to complete dermination was recorded in seeds sown vertically with stalk downward (46.71 days). This could be due to keeping stalk downward needs lot of energy to produce plumule and lack of aeration. The results are in conformity with Kevin et al. (2015) who reported a significant difference in germination speed of Lagenaria siceraria sown in a different orientation.

The interaction between potting mixture and sowing orientation was also noticed significant difference in time take to complete germination. Significantly, minimum number of days taken to complete germination were recorded in potting mixture Soil:Sand:FYM with seeds sown vertically with stalk upward (18.00 days). However, significantly maximum number of days taken to complete the germination were recorded in potting mixtureSoil: Paddy husk : FYM (2:1:1) with seeds sown vertically with stalk downwards (49.33 days). Similar results were also reported by Lucky & Usman (2022).

# 3.3 Seedling Length

The study on seedling growth was also significantly differed with potting mixture and seed orientation. Significant differences were observed among the different treatments with regard to seedling length was observed in seed sown horizontally in potting mixture Soil: Cocopeat: Vermicompost (2:1:1) and results presented in Table 4.

Germination percentage													
		2	022			2	023			Pooled			
	S <sub>1</sub> S <sub>2</sub> S <sub>3</sub> Mean				S₁	S <sub>2</sub>	S₃	Mean	S₁	S <sub>2</sub>	S₃	Mean	
M <sub>1</sub>	42.33	80.33	83.67	68.78	41.00	81.33	85.00	69.11	41.67	81.00	84.33	69.00	
M <sub>2</sub>	35.00	68.00	71.67	58.22	32.33	69.00	73.67	58.33	33.67	68.67	72.67	58.33	
Мз	33.67	71.00	74.33	59.67	30.33	72.33	74.67	59.11	32.33	71.67	74.67	59.56	
M4	25.33	47.33	56.00	42.89	21.00	46.00	54.33	40.44	23.33	47.00	55.33	41.89	
M5	27.67	49.67	59.67	45.67	26.67	51.33	57.67	45.22	27.67	50.67	58.67	45.67	
M6	28.67	54.33	62.67	48.56	27.00	48.33	63.67	46.33	28.00	51.67	63.33	47.67	
M7	31.67	57.33	65.33	51.44	31.66	57.33	58.33	49.11	31.67	57.33	62.33	50.44	
	32.05	61.14	67.62		30.00	60.81	66.76		31.19	61.14	67.33		
	S.Em.±	CD @ 1 %		S.Em.±	CD @ 1	CD @ 1 %			CD @ 1 %				
Factor A	0.452	1.295			0.738	2.114			0.499	1.428			
Factor B	0.296	0.848			0.483	1.384			0.356	0.935			
ΑΧΒ	0.783	2.244			1.279	3.662			0.864	2.474			

Table 2. Effect of planting mixture and seed sowing orientation on seed germination in balanites roxburghii

M1: Soil:Sand: FYM 2:1:1

M3: Soil: Coco peat: Vermicompost 2:1:1

M5: Soil : Paddy husk : Vermicompost : 2:1:1

M7: Soil : saw dust : vermicompost 2:1:1

S1 : Seed laid vertically with stalk end downwards

S3 : Seeds laid vertically with stalk end upwards

M2: Soil: Coco peat: FYM 2:1:1 M4: Soil: Paddy husk : FYM : 2:1:1 M6: Soil : Saw dust : FYM 2:1:1 S2 : Seeds laid horizontally

487

Days taken for germination												
		2	022		-	2	2023			Pooled		
	S <sub>1</sub>	$S_1$ $S_2$ $S_3$ Mean				S <sub>2</sub>	S <sub>3</sub>	Mean	S₁	S <sub>2</sub>	S <sub>3</sub>	Mean
M <sub>1</sub>	44.00	25.67	18.33	29.33	43.33	22.33	17.67	27.78	44.00	24.33	18.00	28.78
M <sub>2</sub>	47.67	33.67	22.33	34.56	48.00	31.67	22.00	33.89	48.00	33.00	22.33	34.44
Mз	45.33	38.33	27.67	37.11	47.00	35.33	29.67	37.33	46.67	37.00	28.67	37.44
M4	49.67	43.33	39.00	44.00	48.33	41.67	39.00	43.00	49.33	42.67	39.33	43.78
M <sub>5</sub>	47.00	47.33	40.67	45.00	47.67	49.33	40.33	45.78	47.67	48.67	40.67	45.67
M <sub>6</sub>	45.33	41.67	39.33	42.11	48.33	43.67	37.00	43.00	47.00	43.00	38.33	42.78
M <sub>7</sub>	42.67	35.67	32.00	36.78	45.67	36.00	32.00	37.89	44.33	36.00	32.00	37.44
	45.95	37.95	31.33		46.91	37.14	31.10		46.71	37.81	31.33	
	S.Em.±	CD @ 1	CD @ 1 %		S.Em.±	CD @ 1	CD @ 1 %		S.Em.±	CD @ 1 %		
Factor A	0.396	1.135			0.806	2.307			0.544	1.559		
Factor B	0.259	0.743			0.527	1.511			0.356	1.021		
A XB	0.686	1.965			1.395	3.996			0.943	2.700		

Table 3. Effect of planting mixture and seed sowing orientation on days taken for germination in balanites roxburghii

M1: Soil:Sand: FYM 2:1:1

M3: Soil: Coco peat: Vermicompost 2:1:1

M5: Soil : Paddy husk : Vermicompost : 2:1:1

M7: Soil : saw dust : vermicompost 2:1:1

S1 : Seed laid vertically with stalk end downwards

M2: Soil: Coco peat: FYM 2:1:1

M4: Soil: Paddy husk : FYM : 2:1:1

M6: Soil : Saw dust : FYM 2:1:1

S2 : Seeds laid horizontally

S3 : Seeds laid vertically with stalk end upwards

Seedling length (cm)													
			2022				2023		Pooled				
	$S_1$ $S_2$ $S_3$ Mean $S_1$					S <sub>2</sub>	S <sub>3</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	S <sub>3</sub>	Mean	
M <sub>1</sub>	21.00	49.67	40.67	37.11	19.67	55.00	43.00	39.22	20.67	52.67	42.00	38.44	
M <sub>2</sub>	35.67	51.33	45.33	44.11	37.33	50.67	46.00	44.67	37.00	51.33	46.00	44.78	
Mз	43.67	64.33	55.33	54.44	41.67	63.00	52.33	52.33	42.67	64.00	54.33	53.67	
M4	22.33	40.33	31.33	31.33	20.00	42.33	33.33	31.89	21.33	41.67	32.33	31.78	
M <sub>5</sub>	26.67	44.67	33.67	35.00	29.00	43.33	36.33	36.22	28.00	44.33	35.33	35.89	
M <sub>6</sub>	24.33	42.33	34.00	33.56	25.67	37.00	30.33	31.00	25.33	39.67	32.33	32.44	
M7	26.67	45.00	35.00	35.56	26.67	45.00	35.00	35.56	26.67	45.00	35.00	35.56	
	28.62	48.24	39.33		19.67	55.00	43.00	39.22	28.81	48.38	39.62		
	S.Em.±	CD @ 1 %			S.Em.±	CD @ 1	CD @ 1 %			CD @ 1 %			
Factor A	0.741	2.121			0.773	2.115			0.571	1.636			
Factor B	0.485	1.389			0.506	1.450			0.374	1.071			
A XB	1.283	3.674			1.339	3.836			989	2.834			

#### Table 4. Effect of planting mixture and seed sowing orientation on seedling length in balanites roxburghii

M1: Soil:Sand: FYM 2:1:1

M3: Soil: Coco peat: Vermicompost 2:1:1

M5: Soil : Paddy husk : Vermicompost : 2:1:1

M7: Soil : saw dust : vermicompost 2:1:1

S1 : Seed laid vertically with stalk end downwards

M2: Soil: Coco peat: FYM 2:1:1

M4: Soil: Paddy husk : FYM : 2:1:1

M6: Soil : Saw dust : FYM 2:1:1

S2 : Seeds laid horizontally

S3 : Seeds laid vertically with stalk end upwards

Seedling vigour Index														
		202	22			20	2023				Pooled			
	S₁	S <sub>2</sub>	S₃	Mean	S <sub>1</sub>	S <sub>2</sub>	S₃	Mean	S₁	S <sub>2</sub>	S₃	Mean		
<b>M</b> 1	888	3989	3403	2760	807	4471	3656	2978	848	4230	3529	2869		
M2	1248	3493	3247	2663	1209	3500	3392	2700	1229	3497	3320	2682		
Мз	1470	4568	4113	3384	1263	4554	3903	3240	1367	4561	4008	3312		
M4	567	1910	1754	1410	418	1948	1807	1391	493	1929	1781	1401		
M <sub>5</sub>	738	2215	2005	1653	778	2227	2096	1700	758	2221	2051	1677		
M <sub>6</sub>	695	2300	2132	1709	694	1785	1936	1472	695	2043	2034	1591		
M7	844	2579	2286	1903	844	2579	2000	1808	844	2579	2143	1855		
	921	3008	2706		859	3009	2684		890	3009	2695			
	S.Em.±	CD @ 1 %		S.Em.±	CD @ 1 %			S.Em.±	CD @ 1 %					
Factor A	43.534	124.684			152.894	152.894			35.684	.684 102.200				
Factor B	28.500	81.625			34.948	100.093			23.361	66.906				
ΑΧΒ	75.404	219.959			92.464	92.464			61.807	177.016	;			

Table 5. Effect of planting mixture and seed sowing orientation on seedling vigour index in balanites roxburghii

M1: Soil:Sand: FYM 2:1:1

M3: Soil: Coco peat: Vermicompost 2:1:1

M5: Soil : Paddy husk : Vermicompost : 2:1:1

M7: Soil : saw dust : vermicompost 2:1:1

S1 : Seed laid vertically with stalk end downwards

M2: Soil: Coco peat: FYM 2:1:1

M4: Soil: Paddy husk : FYM : 2:1:1

M6: Soil : Saw dust : FYM 2:1:1

S2 : Seeds laid horizontally

S3 : Seeds laid vertically with stalk end upwards



Plate 1. Effect of sowing orientation on seed germination and seedling growth

The highest seedling length (64.00 cm) was recorded in seed sown horizontally in potting mixture Soil: Cocopeat: Vermicompost (2:1:1), where as the lowest seedling length (20.67 cm) was recorded in potting mixture Soil:Sand:FYM (2:1:1) with seed sown vertically stalk end downwards. Vermicompost provides adequate nutrients and enhances both the physical properties and the water holding capacity (Soegiman, 1982). The application of cocopeat and vermicompost showed significant effect on seedling length probably due to the synergistic combination of both factors in improving the physical conditions of the media and nutritional factors (Sahni et al., 2008). However, the air filled porosity (AFP), easily available water (EAW) and aeration of vermicompost and FYM were not at the recommended level which in turn limit the root growth and lowered the water holding capacity. Therefore, the potting mixture along with soil, vermicompost and cocopeat is more suitable than vermicompost alone because of the better physical properties and enhanced nutrient level.

# 3.4 Seedling Vigour Index

Seedling vigour index was affected significantly by potting mixture and seed sown orientation (Table 5). Seedling vigour index is depends on seed germination and seedling length. Even though highest germination (84.00%) recorded in Soil:Sand:FYM but this media not provide the nutrients for seedling development since poor nutrient supply by this media, but potting mixture Soil:cocopeat:vermicompost provide more nutrients for seedling development in early stages recorded high seedling vigour index (4561) and high vigour seedlings may survive even in unfavorable conditions and survival rate is also high compare to others.

The probable reasons for the best performance of soil, cocopeat and vermicompost are high organic matter content which increases the water and nutrient holding capacity of the medium, which improve the water utilization capacity of plant. Vermicompost is reported to have bioactive principles which are considered to be beneficial for root growth and greater root initiation, higher germination, increased biomass, enhanced growth and development (Bachman & Metzger, 2008) and also balanced composition of nutrients present in the media (Zaller, 2007). The higher available well decomposed organic matter (Vermicompost) may preserve soil humidity, increase nutrient content and improve soil structure which increase water absorption and maintains the cell turgidity, cell elongation and increase respiration at optimum level, leading to favourable seed sprouting. Vermicompost mixed with soil affects properties of soil physics, chemistry and biology, since organic matter acts as glue for soil aggregate and source of soil nutrient (Soepardi, 1983). Vermicompost develop aggregate. granules may soil Vermicompost and soilmay decrease fluctuation of soil temperature. Further, seed germination and root growth becomes easier to the particular depth so that plant grows well and may absorb

more water and nutrient. Organic matter may also improve nutrient availability and improve phosphorous absorption (Karama and Manwan, 1990). All these factors are favourable for seed germination and ultimate by increasing seed germination percent, seedling length, seedling vigour index and minimizing the days taken for initiation of germination. Combined application of soil, vermicompost and cocopeat in the treatment T3 showed significant effect on germination, seedling growth and vigour of seedling. This may be due to the synergistic combination of both factors in improving physical condition of the media and nutritional factors (Sahni et al., 2008) and Sajana et al. (2018).

# 4. CONCLUSION

Based on the results, it can be concluded that depulped seeds sown in the potting media of Soil:Sand:FYM (2:1:1), with seeds laid vertically (stalk end upwards), yield good germination. However, seedlings from this mix should be transplanted to a different potting medium for better growth and development. Alternatively, when depulped seeds are sown in a mix of Soil:Cocopeat:Vermicompost (2:1:1) with the seeds laid horizontally, there is no need to transplant them to another medium until they are transplanted to the main field because cocopeat and vermicompost provide nutrients to seedlings method allows for initially. This better establishment of the seedlings.

# DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

# COMPETING INTERESTS

Authors have declared that no competing interests exist.

# REFERENCES

- Abdul-Baki, A. A., & Anderson, J. D. (1973). Vigour determination in soybean by multiple criteria. *Crop Science, 13*, 630– 633.
- Ahmed, A. I., Aref, M. I., El-Atta, M. I., & Iqbal, M.
  I. (2017). Enhancing propagation and growth of *Balanites aegyptiaca* through

seed treatment. *Journal of Experimental Biology*, *38*(4), 617–622.

- Akanbi, W. B., Togun, A. O., Adediran, T. A., Olabde, O. S., & Olanivi, J. O. (2002). Effect of split application of organomineral fertilizer on okra growth, nutrient uptake, and fruit yield. *Crop Research*, 29(2), 223– 239.
- Al-Thobaiti, S. A., & Zeid, I. M. A. (2018). Medicinal properties of desert date plants (*Balanites aegyptiaca*) – An overview. *Global Journal of Pharmacology, 12*(1), 1– 12.
- Bachman, G. R., & Metzger, J. D. (2008). Growth of bedding plants in commercial potting mixture substrate amended with vermicompost. *Bioresource Technology*, *99*, 3155–3161.
- Bowers, S. A., & Hayden, C. W. (1972). Influence of seed orientation on bean seedling emergence. *Agronomy Journal*, *64*, 736– 738.
- Chapagain, B. P., Yehoshua, Y., & Wiesman, Z. (2009). Desert date (*Balanites aegyptiaca*) as an arid lands sustainable bioresource for biodiesel. *Bioresource Technology, 100*, 1221–1226.
- Chhanval, A. R., Jadhav, A. L., Jarhad, A. P., Gavhane, V. G., & Krushna, J. (2023). A review on pharmacological activity and interaction of *Balanites roxburghii* (desert date). *International Journal of Novel Research and Development*, 8(5), 84–89.
- Daya, L. C., & Vaghasiya, H. U. (2011). A review on *Balanites aegyptiaca* Del (desert date): Phytochemical constituent, traditional uses, and pharmacological activity. *Pharmacogen Review*, *5*(9), 55–62.
- Elffel, A. A. (2012). Effect of seed pretreatment and sowing orientation on germination of *Balanites aegyptiaca* (L.) Del. seeds. *American-Eurasian Journal of Agricultural* & Environmental Sciences, 12, 897–900.
- El-Nour, M., & Kalislo, M. (1995). Effect of pulp and positioning of seeds on germination and juvenile development of heglig (*Balanites aegyptiaca* (L.) Del.). *Journal of Agricultural Sciences, University of Khartoum, 3*(1), 87–97.
- Hammiche, H., & Maiza, K. (2006). Traditional medicine of leaves in Sahara. Pharmacopocia of Tassili. *Journal of Ethnopharmacology, 105*.
- Karama, A. S., & Manwan, I. (1990). Penggunaan pupuk organik pada tanaman pangan. Makalah pada Lokakarya

Nasional Efisiensi Penggunaan Pupuk. Cisarua Bogor.

- Kevin, K. K., Bernard, N. K., Laurent, K. K., Ignace, K. K., Pierre, B. J., & Arsène, Z. B. (2015). Effects of seed orientation and sowing depths on germination, seedling vigor, and yield in oleaginous type of bottle gourd, Lagenaria siceraria (Molina Standl). International Research Journal of Biological Sciences, 4(12), 46–53.
- Orwa, C., Mutua, A., Kindt, R., Jamnadass, R., & S., A. (2009). Agroforestry database: A tree species reference and selection guide version 4.0. World Agroforestry Centre (ICRAF), Nairobi, Kenya.
- Saboo, S. S., Chavan, R. W., Tapadiya, G. G., & Khadabadi, S. S. (2014). An important ethnomedicinal plant *Balanites aegyptiaca* Del. *International Journal of Phytopharmacy*, *4*(3), 75–78.
- Sahni, S., Sharma, B. K., Singh, D. P., Singh, H. B., & Singh, K. P. (2008). Vermicompost enhances performance of plant growthpromoting rhizobacteria in *Cicer arietinum*

rhizosphere against Sclerotium rolfsii. Crop Protection, 27, 369–376.

- Sajana, S., Munde, G. R., & Shirsath, A. H. (2018). Effect of growing media on seed germination and seedling growth of marking nut (*Semecarpus anacardium*). *Plant Archives*, 18(SI), 19–26.
- Soegiman. (1982). Ilmu tanah. Terjemahan dari *The nature and properties of soils*. Buckman and Brady. Bhatara Karya Aksara. Jakarta. 788 hal.
- Soepardi, G. (1983). Sifat dan ciri tanah. Department Ilmu-Ilmu Tanah, IPB, Bogor.
- Wakawa, L. D., & Suleiman, U. M. (2022). Germination characteristics of *Balanites* aegyptiaca (L.) Del. seeds under varying light intensity and sowing orientations in a Sudano-Sahelian zone of Nigeria. *Eurasian Journal of Forest Science*, 10(3), 64–71.
- Zaller, J. G. (2007). Vermicompost as a substitute for peat in potting media: Effects on germination, biomass allocation, yields, and fruit quality of three tomato varieties. *Scientia Horticulturae*, *112*, 191–199.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/126176