

Journal of Advances in Biology & Biotechnology

Volume 27, Issue 11, Page 1555-1562, 2024; Article no.JABB.127543 ISSN: 2394-1081

Quality Evaluation of Green Banana (cv. Nendran) in Different Stages of Maturity

Shihna a++ and Suma Divakar a#*

^a Department of Community Science, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala Agricultural University, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/jabb/2024/v27i111742

Open Peer Review History: This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/127543

Original Research Article

Received: 10/11/2024 Accepted: 02/12/2024 Published: 05/12/2024

ABSTRACT

Aims: Banana (*Musa spp.*) is a globally significant tropical fruit crop, valued for its economic importance and as a staple food that contributes significantly to human nutrition. The Nendran banana variety (Musa AAB) is widely cultivated in the South Indian states of Tamil Nadu, Kerala, and Karnataka, and is also well-known in other regions worldwide. This study aims to examine the physical, chemical, and nutritional characteristics of Nendran bananas at various maturity stages—75, 80, 85, and 90 days after flower emergence.

Study Design: Completely randomized design (CRD).

Place and Duration of Study: Department of Community Science, College of Agriculture, Vellayani, Kerala Agricultural University, between February 2024 to October 2024.

++ M.Sc. Scholar;

Professor;

*Corresponding author: E-mail: suma.divakar@kau.in;

Cite as: Shihna, and Suma Divakar. 2024. "Quality Evaluation of Green Banana (cv. Nendran) in Different Stages of Maturity". Journal of Advances in Biology & Biotechnology 27 (11):1555-62. https://doi.org/10.9734/jabb/2024/v27i111742.

Methodology: Nendran bananas for this study were obtained from a local farmer in Vellayani, Thiruvananthapuram. Maturity was monitored by tagging the plants on the day of flower emergence, and the fruits were harvested at intervals of 75,80,85, and 90 days post-emergence. At each stage, green fruits were selected from the middle three fruits of the third hand from the top of each bunch and taken for physical (fruit weight, fruit length, circumference, peel weight, pulp weight), chemical (moisture, acidity, TSS, crude fiber and tannin) and nutritional (starch, protein, total sugar, reducing sugar) analysis.

Results: The result indicate that the ripening process significantly impacts the physical, chemical, and nutritional properties of Nendran bananas. Physical attributes such as fruit weight, pulp weight, fruit length, and diameter reached their maximum at 90 days, while peel weight was highest at 75 days. Chemical analysis demonstrated that moisture content, acidity, and total soluble solids (TSS) were greatest at 90 days, whereas crude fiber and tannin content peaked at 75 days. Nutritional analysis revealed that protein, total sugar, and reducing sugar levels were highest at 90 days, while starch content was highest at 75 days. These findings highlight how understanding the nutritional profile of unripe green Nendran bananas at different maturity stages offers valuable insights into the chemical transformations that occur throughout the ripening process.

Keywords: "Nendran"; maturity; green banana; nutrients; physico-chemical.

1. INTRODUCTION

Banana (*Musa spp.*) is a major tropical fruit crop holdina substantial alobally. economic importance and serving as a staple food with a vital role in human nutrition (Aurore et al., 2009). India is the world's top banana grower, with China, Ecuador, Brazil, and the Philippines following closely after (Nayar, 2010). Bananas are nutritious, sweet, and readily digestible fruits high in carbohydrates and minerals such as potassium, magnesium, salt, and phosphorus. They even have more calories than potatoes (Pereira and Maraschin, 2015). Bananas are unique because they combine energy-boosting components with tissue-building nutrients. including proteins, vitamins, and minerals. (Qamar and Shaikh, 2018). They can contribute significantly to the vitamin A, C, and B6 levels in the diet while also providing an instant source of energy (Singh et al., 2016).

The raw banana variety Musa (AAB) cv. "Nendran" is extensively cultivated in the South Indian states of Tamil Nadu, Kerala, and Karnataka, and is also popular in other regions worldwide. Kerala now generates less than 30% of its banana requirements and relies on Tamil Nadu to assist cover the gap (Hassan, 2016). The state is home to a wide assortment of banana cultivars, each having its own regional preferences and economic worth. Among them, the "Nendran" variety is very popular in Kerala and is widely planted for both fresh fruit consumption and banana chip production. It is enjoyed in both unripe and ripe stages. "Nendran" belongs to the French plantain family and is an important part of the local diet. However, the area allocated to banana production in Kerala is shrinking as farmers turn to more profitable crops. Nonetheless, bananas' profound historical, economic, and even religious significance in Kerala helps to keep the crop from being fully eclipsed by other agricultural endeavors (Siji and Nandini, 2017).

"Nendran" bananas mature 5-6 months after blossoming, producing large, thick fruits with a high shelf life. Unripe "Nendran" is ideal for Kerala banana chips and is often used as a first weaning snack for infants in India (Chitra, 2015). The fruits average 20-27 cm in length, 14-18 cm in diameter, and have a thick peel with three ridges and a unique tip. A Nendran bunch weighs 12-15 kg, consisting of 4-6 hands with 8-10 fingers each. The creamy pale-yellow flesh of unripe fruits deepens to vellowish-orange as they ripen, while the peel transitions from green to vellow and later develops dark spots (Sreejith and Sabu, 2017). Unripe flesh has a bland, starchy texture, which becomes sweet and creamy as starch converts to sugar (Suman et al., 2018). "Nendran" flower bracts are rich in anthocyanins with anticancer properties, its peel contains a metalloprotease with collagenolytic and cytotoxic effects, and its fructans act as prebiotics by promoting lactic acid bacteria (Shalini et al., 2017).

2. MATERIALS AND METHODS

2.1 Collection of Raw Materials

In this study, "Nendran" bananas were sourced from a local farmer at Vellayani,

Thiruvananathapuram. To accurately determine maturity, the bananas were tagged on the day of flower emergence. Fruits were harvested at 75, 80, 85,and 90 days post-emergence. For each maturity stage, green fruits were collected from the middle three fruits on the third hand from the top of each bunch and brought to the laboratory for subsequent physical and biochemical analyses.

2.2 Physical Parameters

The physical parameters of unripe green banana fruits were assessed using the method described by Dadzie and Orchard (1997).

The length of each fruit was measured along its outer curve from the distal end using a measuring tape. Measurements were taken from a minimum of five fruits. The circumference of each fruit was measured at its widest midpoint using a measuring tape, ensuring consistency in the measurement process. Readings were collected from three fruits at each maturity stage to provide a representative assessment of fruit circumference across different developmental phases. In addition, the weight of the unripe green fruits, as well as the individual weights of the pulp and peel, were accurately determined using an electronic balance. This method allowed for precise quantification of the various components of the fruit, facilitating detailed analysis of their physical characteristics.

2.3 Chemical Parameters

For the analysis of chemical parameters, the fruit pulp was cut into uniform pieces and immersed in a pretreatment solution (Karthikeyan, 2015) to minimize browning reactions. Subsequently, the treated pulp was ground into a fine paste for further analysis.

2.3.1 Moisture content

Moisture content was determined according to the AOAC (2010).

2.3.2 Titratable acidity

Titratable acidity was measured according to the method established by Ranganna, (1977). A known volume of the fruit sample was taken and titrated with a 0.1 N sodium hydroxide (NaOH) solution, using phenolphthalein as an indicator. The acidity levels were reported as a percentage (%).

2.3.3 Total soluble sugars

Total soluble solids (TSS) were measured using a hand refractometer with a range of 0 to 32 °Brix and were expressed in degrees Brix (°Brix) according to the method described by Ranganna (1977).

2.3.4 Crude fiber

Crude fiber content was determined using the acid-alkali digestion method as recommended by Chopra and Kanwar (1978).

2.4 Nutritional Parameters

2.4.1 Starch content

Starch content was determined using the AOAC(2010) method.

2.4.2 Protein

The protein content in banana samples was determined using the micro Kjeldahl wet digestion method (AOAC,2010).

2.4.3 Total sugar

Total sugars in the sample were estimated following the AOAC (2010) method and expressed as a percentage on a fresh weight basis.

2.4.4 Reducing sugar

Reducing sugars were determined by the copper reduction method using Fehling's solution (AOAC,2010) and expressed as a percentage.

3. RESULTS AND DISCUSSION

Significant changes were observed in the physical, chemical, and nutritional parameters of bananas at different maturity stages. The variation in these parameters highlights how maturation impacts the fruit's composition, including aspects such as texture, sugar content, acidity, and nutrient levels. This underscores the importance of maturity in determining the quality and nutritional profile of bananas, influencing both their consumer acceptability and potential applications in food processing and nutrition.

3.1 Physical Parameters

The physical characteristics of banana fingers showed significant variations across the different

maturity stages tested (Table 1). Fruit weight was notably affected by the stage of maturity, with the 90-day maturity stage (149.25) exhibiting the highest fruit weight, while the 75-day stage recorded the lowest among those tested. In terms of peel and pulp weight, the 75-day maturity stage had a higher peel weight but a lower pulp weight. Conversely, the 90-day maturity stage showed an increased pulp weight, highlighting the shift in fruit composition with advancing maturity. Fruit length was also influenced by maturity stage, with the 90-day maturity showing the greatest length compared to the other stages. The 75-day and 80-day maturity stages exhibited similar fruit lengths, indicating minimal variation between them. In terms of fruit diameter, the 90-day maturity stage recorded the highest diameter, while the other maturity stages showed similar measurements. Comparable changes have been documented by Rodriguez-Sosa et al. (1977) in their research on banana flour preparation.

3.2 Chemical Composition of Banana

The chemical composition of green Nendran bananas were evaluated, showing significant differences across various maturity stages (Table 2). This analysis highlights the unique chemical composition at each stage, demonstrating the progression of changes as the fruit matures. These findings underscore the importance of understanding such differences for applications in nutritional profiling and postharvest management.

The moisture content in bananas is observed to be highest at 90 days of maturity (59.410%) and lowest at 75 days (49.603%). Typically, as bananas undergo the ripening process, their moisture content ranges from 60% to 68.6%, and progressively increases to levels between 68.6% and 78.1% (Sakyi et al., 2008). This increase in moisture content during ripening is likely attributed to the breakdown of carbohydrates, which leads to the softening of tissue texture, as well as osmotic transfer from the peel to the pulp (Onwuka, G.I. and Onwuka, N.D. 2005).

The acidity of banana fingers has been analyzed, and findings indicate that the highest acidity levels are observed at 90 days of maturity compared to other stages. This suggests that as bananas progress through their ripening process, the acidity initially increases and peaks at the later stage of 90 days. High acidity at this point is often associated with the biochemical changes that occur as the fruit ripens, including the conversion of starches into sugars and the formation of organic acids (wyman, 1964). These changes contribute to the flavor profile of the banana, enhancing its overall taste and sensory characteristics, Sreedevi and Suma (2015) reported similar results, as did Mayadevi (2016).

| Maturity (Days) | Fruit weight (g) | Peel weight (g) | Pulp weight (g) | Fruit length (cm) | Fruit diameter (cm) |
|--------------------|---------------------|--------------------|---------------------|----------------------|------------------------|
| 75 | 115.25 ^b | 50.75 ^a | 56.50 ° | 18.175 ^b | 11.175 ^b |
| 80 | 109.50 ^b | 39.50 bc | 66.25 ^b | 18.350 ^b | 11.350 ^b |
| 85 | 116.50 ^b | 39.0 ^c | 75.50 ^{ab} | 21.05 ^{ab} | 11.625 ^{ab} |
| 90 | 149.25 ^a | 48.0 ^{ab} | 81.25 ^a | 22.325 ^a | 12.275 ^a |
| SE (d) | 8.215 | 3.999 | 4.26 | 1.351 | 0.316 |
| SE(m) | 5.809 | 2.828 | 3.012 | 0.955 | 0.223 |
| CD | 17.898 | 8.712 | 9.281 | 2.943 | 0.688 |

Table 1. Physical parameters of banana fingers

Table 2. Chemical parameters of banana fingers

| SI.no | Maturity (Days) | Moisture (%) | Acidity (%) | TSS (º brix) | Crude fiber (%) | Tannin (g 100 g⁻¹) |
|-------|--------------------|---------------------|--------------------|---------------------|--------------------|-----------------------|
| 1 | 75 | 49.603 ^d | 0.133 ^d | 4.800 d | 4.043 a | 8.017 ^a |
| 2 | 80 | 53.457 ° | 0.160 ° | 7.013 ° | 3.337 ^b | 7.347 ^b |
| 3 | 85 | 55.813 ^b | 0.190 ^b | 10.900 ^b | 3.290 ° | 6.770 ° |
| 4 | 90 | 59.410 ^a | 0.233 ^a | 14.200 ^a | 2.850 d | 5.897 ^d |
| | SE (d) | 0.026 | 0.008 | 0.002 | 0.008 | 0.009 |
| | SE(m) | 0.018 | 0.006 | 0.002 | 0.006 | 0.006 |
| | CD | 0.06 | 0.018 | 0.005 | 0.018 | 0.021 |

| SI. no | Maturity (Days) | Starch (mg) | Protein (mg) | Total sugars (%) | Reducing sugars (%) |
|--------|--------------------|-------------------|-------------------|---------------------|------------------------|
| 1 | 75 | 84.0 ª | 1.50 d | 1.83 ^d | 0.250 d |
| 2 | 80 | 83.5 ^b | 1.90 ° | 2.10 ° | 0.450 ° |
| 3 | 85 | 80.1 ° | 2.20 ^b | 2.610 ^b | 0.806 ^b |
| 4 | 90 | 76.4 ^d | 2.31 ^a | 2.806 ^a | 1.033 ^a |
| | SE (d) | 0.071 | 0 | 0.025 | 0.024 |
| | SE(m) | 0.05 | 0 | 0.018 | 0.017 |
| | CD | 0.163 | 0 | 0.058 | 0.055 |

Table 3. Nutritional parameters of banana fingers

In the analysis of total soluble sugars across different maturity, it was observed that the concentration was highest at day 90 (14.200°brix) and lowest at day 75 (4.800°brix). This trend indicates a significant increase in soluble sugar content as the maturity of the sample advances. The increase in total soluble sugars during ripening is an essential trait that reflects the hydrolysis of starch into soluble sugars, including glucose, sucrose, and fructose (Meghwal et al., 2021). This process underscores the enzymatic conversion that occurs as the fruit matures, contributing to the accumulation of sugars and enhancing the sweetness and overall quality of the produce (Tirkey et al., 2003). The significant rise in total soluble sugars at advanced maturity, such as day 90, highlights the culmination of starch breakdown and sugar synthesis. Converselv, the lower sugar content at day 75 indicates an earlier stage of development where this conversion is less pronounced. These observations are crucial for optimizing harvest timing to achieve the desired sugar profile in agricultural and food production practices (Mahato et al., 2014).

Crude fiber content was found to be highest at the earlier stage of 75 days and decreased significantly by 90 days. This trend was similarly observed in unripe bananas, where the crude fiber composition was initially high but declined as the fruit ripened (Dhaval and Naik, 2010). Barnell HR (1940) noted in his work that the decrease in crude fiber during ripening is due to enzymatic breakdown of the complex into simpler, more easily polysaccharides digestible forms. This transformation, as the fruit ripens from an unripe to a fully mature state, involves the degradation of cell wall components like cellulose and hemicellulose, leading to a softer texture and improved palatability. Understanding this pattern is important for determining optimal harvest times, particularly for applications where fiber content plays a role in the nutritional or functional quality of the produce.

In the present study, tannin content was found to be highest at day 75 and decreased significantly by day 90. This finding aligns with previous research by Khawas et al. (2014), which demonstrated that the tannin content in banana samples varies significantly across different stages of development. The decline in tannin levels during maturation contributes to the sensory evolution of the fruit. As noted by Szajdek and Borowska (2008), tannins play a crucial role in shaping the sensory properties of fruits and fruit products, imparting a characteristic tartness and influencing color changes in both the fruit and its derived juices. The reduction in tannin content at advanced ripening stages corresponds to the softening of the fruit and a shift towards a sweeter taste profile, enhancing consumer appeal and usability in various culinary and processing contexts.

3.3 Nutritional Parameters

The nutritional properties of green Nendran bananas were investigated, with an emphasis on starch, protein, total sugar, and reducing sugars. The research provides a full understanding of the nutritional composition of bananas at various development stages. Starch, being the major carbohydrate in unripe bananas, contributes significantly to the overall energy content and steadily diminishes as the fruit ripens and converts into simpler sugars. Protein levels add to the nutritional value, but in lesser proportions than carbs. Total sugars and reducing sugars, essential for flavor and palatability, were also analyzed, revealing their increase as starch breaks down during the ripening process. Understanding these nutritional profiles is critical for making the best use of green Nendran bananas in dietary planning and food processing applications.

In the current study, the starch content of unripe bananas was shown to fluctuate from 84% at day

75 to 76.4% at day 90. Starch levels peaked at day 75 and dropped as the fruit grew to day 90. According to Butani and Chovatia (2014), the breakdown of starch into simpler sugars during the ripening process results in the sweeter taste and softer texture characteristic of ripe bananas.

In the investigation of green Nendran bananas, the protein content was shown to be greater at day 90 when compared to other maturity stages. This increase in protein content as the fruit matures may be due to biochemical changes that occur during ripening, such as the creation and accumulation of numerous enzymes and structural proteins. While proteins are not the predominant nutritional component of bananas, contribution is considerable their when considering the fruit's total nutritional profile. The rise at day 90 suggests that fully ripened bananas may have more protein content, adding to their nutritional worth (Vanilarasu and This information. Balakrishnamurthy, 2014). together with data on starch conversion and sugar buildup, helps to highlight the overall nutritional advantages of bananas at different stages of maturity (Lohi 2010).

The total sugars in unripe banana pulp were analysed at various development stages. The investigation revealed that the overall sugar level was greatest at day 90 when compared to the other stages. These findings align with previous research by Lodh et al. (1973), which found comparable patterns of total sugar concentration in bananas at various stages of maturation.

Venkatarayappa et al. (1976) and Chellappan (1983) reported that dessert bananas contained significantly higher levels of reducing sugars, whereas cooking bananas and plantains exhibited greater non-reducing sugar content. This study found that reducing sugar levels were highest at the 90 maturity day compared to earlier stages.

4. CONCLUSION

This study reveals how the nutritional makeup of bananas varies with development stage. The data shows that ripening has a substantial influence on the physical, chemical, and nutritional aspects of Nendran bananas. Physical measures such as fruit weight, pulp weight, fruit length, and fruit diameter were higher at day 90, but peel weight was highest at day 75. Chemical analysis revealed that moisture content, acidity, and total soluble solids (TSS) peaked at day 90, whereas crude fibre and tannin levels peaked at day 75. In terms of nutritional characteristics, protein, total sugar, and reducing sugar concentrations were highest at day 90, whereas starch content was highest at day 75. The findings showed that knowing the nutritional quality of unripened green Nendran bananas at various maturity stages provides insights into the chemical changes that occur during ripening. This knowledge can be used to extend the usage of bananas in a variety of culinary items. The outcomes of this study will help the food sector optimize nutritional contents when developing new products that incorporate the examined banana cultivar.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge to the Department of Community Science, College of Agriculture, Vellayani, Thiruvananthapuram for providing the facilities to carry out this research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- AOAC. (2010). Official methods of analysis of the Association of Official Analytical Chemists (18th ed.). AOAC International.
- Aurore, G., Parfait, B., & Fahrasmane, L. (2009). Bananas, raw materials for making processed food products. *Trends in Food Science & Technology, 20*(2), 78–91. https://doi.org/10.1016/j.tifs.2008.10.003
- Barnell, H.R. (1940). Studies in tropical fruits. VII. Carbohydrate metabolism of the banana fruit. *Annals of Botany, 7*(4), 297-309.
- Butani, A.M., & Chovatia, R.S. (2014). Effect of chemical fertilizer and vermicompost on days to flowering, harvesting and maturity of banana (*Musa paradisiaca* L.) cv. Grand Naine. *The Asian Journal of Horticulture*, 9(2), 305–308.

- Chellappan, K. (1983). Effect of 2, 4-D and GA on fruit development and post harvest physiology in bananas. Ph.D. Thesis, Faculty of Horticulture, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.
- Chitra, P. (2015). Development of banana-based weaning food mixes for infants and its nutritional quality evaluation. *Revista de Saúde Ambiental*, *30*(2), 125-130. https://doi.org/10.1515/reveh-2015-0002
- Chopra, S.L., & Kanwar, J.S. (1978). *Analytical agricultural chemistry*. Kalyani Publishers.
- Dadzie, B.K., & Orchard, J.E. (1997). Routine post-harvest screening of banana/plantain hybrids: Criteria and methods (Vol. 2). Bioversity International.
- Dhaval, R.P., & Naik, A.G. (2010). Effect of preharvest treatment of organic manures and inorganic fertilizers on postharvest shelf life of sapota cv. Kalipatti. *Indian Journal of Horticulture, 67*(3), 381–386.
- Hassan, S.N. (2016). A study on technological gap in banana cultivation technologies in southern district of Tamil Nadu. *International Journal of Scientific and Research Publications, 6*(7), 388-394.
- Karthikeyan, N. (2015). Development of value added products from banana peel. M.Sc.(FS&N) thesis, Kerala Agricultural University, Thrissur.
- Khawas, P., Das, A.J., Dash, K.K., & Deka, S.C. (2014). Thin-layer drying characteristics of Kachkal banana peel (*Musa ABB*) of Assam, India. *International Food Research Journal, 21*, 975-982.
- Lodh, S.B., Divakar, N.G., Chadha, K.L., Melanta, K.R., & Selvaraj, Y. (1973). Biochemical changes associated with growth and development of pineapple fruit variety Kew. III. Changes in plant pigments and enzyme activity. *Indian Journal of Horticulture, 30*, 381-383.
- Lohi, D.J. (2010). Medicinal and nutritional values of banana cv. Nendran. *Asian Journal of Horticulture*, *5*(1), 11-14.
- Mahato, S.K., Suresh, C.P., & Kuna, R.K. (2014). Effects of inorganic fertilization on growth, yield, and fruit quality of plantain cv. Nendran. *Plant Archives, 14*(2), 939–944.
- Mayadevi, M.R. (2016). Efficiency of vermiconversion and decomposition of farm residues on soil health, yield, and quality of banana (*Musa* spp.). Ph.D. Thesis, Kerala Agricultural University, 150–198.
- Meghwal, M.L., Jyothi, M.L., Pushpalatha, P.B., & Bhaskar, J. (2021). Yield and fruit quality

of banana *Musa* (AAB) Nendran as influenced by nutrient sources. *International Journal of Bio-resource and Stress Management, 12*(2), 131-136.

- Nayar, N.M. (2010). The bananas: Botany, origin, dispersal. In *Horticultural reviews*. John Wiley & Sons, Inc. https://doi.org/10.1002/9780470527238.ch 2
- Onwuka, G.I., & Onwuka, N.D. (2005). The effects of ripening on the functional properties of plantain and plantain-based cake. *International Journal of Food Properties, 8*(2), 347-353. https://doi.org/10.1081/JFP-200059489
- Pereira, A., & Maraschin, M. (2015). Banana (*Musa* spp) from peel to pulp: Ethnopharmacology, source of bioactive compounds, and its relevance for human health. *Journal of Ethnopharmacology, 160*, 149–163. https://doi.org/10.1016/j.jep.2014.11.008
- Qamar, S., & Shaikh, A. (2018). Therapeutic potentials and compositional changes of valuable compounds from banana – A review. *Trends in Food Science & Technology, 79*, 1–9. https://doi.org/10. 1016/j.tifs.2018.06.016
- Ranganna, S. (1977). *Manual for analysis of fruit and vegetable products*. Tata McGraw-Hill Publication Company Ltd.
- Rodriguez-Sosa, E.J., González, M., Caloni, I., & Parsi-Ros, O. (1977). The preparation of green banana flour. *Journal of Agriculture of the University of Puerto Rico, 61*(4), 470-478.
- Sakyi-Dawson, E., Asamoah-Bonti, P., & Annor, G.A. (2008). Biochemical changes in new plantain and cooking banana hybrids at various stages of ripening. *Journal of the Science of Food and Agriculture, 88*(15), 2724-2729. https://doi.org/10.1002/jsfa. 3399
- Shalini, R., Abinaya, G., Saranya, P., & Antony, U. (2017). Growth of selected probiotic bacterial strains with fructans from Nendran banana and garlic. *LWT - Food Science and Technology*, *83*, 68–78. https://doi.org/10.1016/j.lwt.2017.03.059
- Siji, S., & Nandini, P.V. (2017). Chemical and nutrient composition of selected banana varieties of Kerala. *International Journal of Advanced Engineering, Management and Science, 3*(4), 239829.
- Singh, B., Singh, J.P., Kaur, A., & Singh, N. (2016). Bioactive compounds in banana and their associated health benefits – A

review. *Food Chemistry*, 206, 1–11. https://doi.org/10.1016/j.foodchem.2016.03 .033

- Sreedevi, L., & Suma, D.A. (2015). Comparative quality analysis of banana cv. Palayamkodan. *International Journal of Agricultural Research, 4*(4), 6–11.
- Sreejith, P.E., & Sabu, M. (2017). Edible bananas of South India: Taxonomy & phytochemistry. India: Rheedea.
- Suman, R., Kalaimathi, K., Palanichamy, S., Sowmiya, R., Vaganan, M.M., Ravi, I., et al. (2018). Anti-cancerous activities of anthocyanins of banana cv. Nendran (*Musa* sp.) flower bracts against human colon and cervical cancer cell lines. *International Journal of Current Microbiology and Applied Sciences*, 7(12), 2786–2793.

https://doi.org/10.20546/ijcmas.2018.712.3 16

Szajdek, A., & Borowska, E.J. (2008). Bioactive compounds and health-promoting

properties of berry fruits: A review. *Plant Foods for Human Nutrition, 63,* 147-156.

- Tirkey, T., Agarwal, S., & Pandey, S.D. (2003). Effect of organic manures on growth, maturity, and yield of banana cv. Dwarf Cavendish. *South Indian Horticulture, 50*, 19–24.
- Vanilarasu, G.B., & Soorianatha, S. (2014). Influence of organic manures and amendments on growth, flowering, yield, quality and postharvest life of banana cv. Grand Naine. *International Journal of Environmental Science, 4*, 147–152.
- Venkatarayappa, T., Narasham, B., & Venkatesan, C. (1976). Effect of postshooting application of urea on development and composition of banana fruit. *South Indian Horticulture, 19*, 109–117.
- Wyman, H., & Palmer, J.K. (1964). Organic acids in the ripening banana fruit. *Plant Physiology, 39*(4), 630.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/127543