

Asian Journal of Research in Agriculture and Forestry

Volume 10, Issue 4, Page 514-523, 2024; Article no.AJRAF.128483 ISSN: 2581-7418

Reevaluating Shifting Cultivation in Northeast India: Balancing Traditional Practices, Sustainability, and Development

Ritika Bhardwaj^{a++*}

^a Network for Certification and Conservation of Forests (NCCF), ATS Bouquet, Tower B, B-801(A), 8th Floor, Sector 132, Noida, U.P 201304, India.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: https://doi.org/10.9734/ajraf/2024/v10i4354

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/128483

Review Article

Received: 15/10/2024 Accepted: 19/12/2024 Published: 23/12/2024

ABSTRACT

Shifting cultivation, particularly the jhum system, has been a long-standing practice in Northeast India, sustaining the livelihoods of indigenous communities. This paper aims to reevaluate the enduring practice of shifting cultivation in Northeast India, acknowledging its deep historical roots, vital role in sustaining tribal livelihoods, and dynamic adaptation amidst socioeconomic change. Tracing its origins to the Neolithic era, shifting cultivation has evolved from a subsistence-based model to one generating modest surpluses, remaining integral to food security and cultural identity in rural tribal communities. Various research papers and government reports have been studied thoroughly from 1985 to 2023 to examine the evolving perceptions of shifting cultivation, considering both its economic and environmental implications. While often criticized for contributing

++ Program Officer;

*Corresponding author: E-mail: ritikabhardwaj1575@gmail.com;

Cite as: Bhardwaj, Ritika. 2024. "Reevaluating Shifting Cultivation in Northeast India: Balancing Traditional Practices, Sustainability, and Development". Asian Journal of Research in Agriculture and Forestry 10 (4):514-23. https://doi.org/10.9734/ajraf/2024/v10i4354. to deforestation, soil degradation, and biodiversity loss, recent research suggests that the practice can be more sustainable than traditionally assumed, especially when fallow cycles are maintained. The jhum system promotes biodiversity conservation through its use of mixed cropping and the protection of endemic species. However, challenges such as insecure land tenure and inadequate policies have hindered the adoption of alternative farming practices. This study emphasizes the significance of blending traditional ecological wisdom with contemporary agricultural techniques to bolster food security, conserve biodiversity, and enhance community well-being. The findings demonstrate that, when bolstered by sustainable interventions such as homestead production systems that honor indigenous values, shifting cultivation continues to serve as a crucial livelihood for tribal communities in Northeastern India, fostering rural development, cultural preservation, and environmental sustainability.

Keywords: Shifting cultivation; deforestation; traditional knowledge; biodiversity; fallow cycle.

1. INTRODUCTION

Shifting cultivation has been practiced in the Northeastern Region of India since ancient times, as old as human settlements in the region. It is a traditional agricultural practice carried out across the hilly landscapes of the area [1]. The shifting cultivation, particularly the method of preparing Farm plot, requires clearing a patch of forest by slashing the trees and vegetative materials therein, which are then left to dry before these are set on fire to produce ash for treating the soil and preparing the farm bed for sowing.

Bos et al., 2020 [2], mentions that landscapes influenced by shifting cultivation span approximately 280 million hectares globally, with around 30-40 million individuals relying on this practice in Southeast Asia alone. Shifting cultivation is often portraved as an economically inefficient and environmentally damaging form of agriculture. Although critics argue that the practice harms the environment by degrading the soil, water, and local biodiversity, numerous scientific and agroecological studies have shown this perception to be inaccurate [3], various scientific and agroecological studies prove otherwise. Terming this view as a 'false narrative', proponents of the practice say that shifting cultivation is perhaps more sustainable settled agriculture and monoculture; than practices that the government overtly promotes in the region [4,5,6].

The jhum system is a complex practice that interacts with both socio-economic and environmental factors. It is based on principles of common property, self-governance, shared labour, and traditional ecological knowledge. Additionally, it maintains a subsistence-based economy and a food production system that safeguards and sustains its own seed varieties and cultivation practices, which have been orally transmitted across generations for centuries. Many reports have highlighted the environmental concerns associated with shifting cultivation. including soil erosion, degradation, and loss of arable land, as well as biodiversity loss, deforestation, and carbon emissions, all of which contribute to ecological imbalances and climate change [7]. Jhum is distinguished by mixed cropping, which mimics nature regarding species diversity and is vital in the in-situ preservation of numerous varieties of edible food crops. Despite facing various criticisms, the jhum landscapes is believed to serve as repositories for the original gene pools of biodiversity, which are conserved and maintained through indiaenous food practices the ingenious production and management of fallow forests in the hills. While some modern economists and environmentalists perceive shifting cultivation as detrimental to the ecosystem, portraying it as disadvantageous and exerting negative impacts on the economy [8], the others maintain that practitioners of this practice are integral to conservation efforts [9,10]. Jhum is distinguished by mixed cropping, which mimics nature regarding species diversity and is vital in the in-situ preservation of numerous varieties of edible food crops [6]. Shifting cultivation has been practiced for generations, encompassing ancestral wisdom in food production, seed conservation, and farm management. This agricultural system has demonstrated remarkable resilience and adaptability, particularly in terms of crop selection and soil nutrient management. However, the primary concern is that the reduced fallow period is not allowing sufficient time for the land to naturally regenerate and restore its soil health.

Bawa et al.,1997 [11] further emphasized that deforestation and habitat fragmentation present more significant threats to biodiversity and wild

genetic resources compared to climate change. Thus, whether we need to go for improved production or conserve the existing rich diversity of crops cultivated in shifting cultivation systems remains a dilemma that requires careful consideration [12]. Various efforts have been made till date to adapt to different strategies by various institutes such as ICAR, ICIMOD, Wildlife Works Carbon (WWC) and many other in different states of the region had met with limited success. This can be attributed to factors such as complex land tenure arrangements and inadequate infrastructure for marketing pose significant challenges. Furthermore, the introduction of modern agroforestry or the complete eradication of traditional shifting cultivation and associated local knowledge to promote monoculture cash crops has not been effectively integrated with meaningful economic improvements for the indigenous communities. This lack of a holistic approach that respects the socio-cultural context and traditional livelihood strategies has hindered the successful adoption of alternative practices in the region.

2. METHODOLOGY

The present paper is an attempt to analyze the practice of shifting cultivation in Northeastern region as an economic activity in the light of the changing perceptions on shifting cultivation in the region. The study is descriptive in its design and has utilized a qualitative approach. The paper mostly reviews the secondary literature hitherto available on shifting cultivation in the region and synthesizes the implications with respect to the changing perceptions during the last three decades (1985-2023). This paper argues that eradicating shifting cultivation is not a viable solution, but rather the solution lies in building upon and improving this traditional practice. The paper emphasizes that adopting alternatives such as cash crop monoculture is not an effective solution for protecting the interests of indigenous communities and the environment. It advocates for a more nuanced approach that recognizes the value of shifting cultivation and explores ways to enhance its sustainability and benefits to local populations and ecosystems.

2.1 Environmental Impacts of Shifting Cultivation and the Role of Fallow Cycles

The jhum cultivation system practiced by upland communities in India's northeastern region utilizes the ecological resources of forests to maintain soil health and enable food production on hillsides. However, concerns have been

raised about the environmental impact of this practice. While shifting agriculture is commonly linked to low crop yields and rapid soil deterioration, emerging research suggests the traditional view of shifting cultivation as environmentally detrimental in terms of its impacts on soil is not well established [13]. A study conducted by Mukul et al.,2022 [14] found that shifting cultivation may not be as detrimental to soil quality, at least in the specific soil type and climate examined. The results indicate that geographic and site-specific factors can play a crucial role in shaping the impact of shifting cultivation on soil properties. The analysis suggests that patch size, slope, and fallow duration are important variables that influence the recovery process of soil macronutrients [15].

According to traditional knowledge jhum creates secondary vegetation that is preferred by herbivores like elephants, sambar and gaur and thus help to coexist. However, Where the jhum cycle has been reduced to 2-3 years, this may have detrimental impacts on the ecosystem and wildlife habitats. Consequently, reducing fallow periods is a concerning prospect. A long fallow period is crucial for restoring essential soil nutrients, and it may require up to 20 years to rehabilitate soil nutrient levels comparable to undisturbed forests [16, 17]. Research conducted by Saxena et al., 1984 shows that fallow cycles as short as 4-5 years in northeast India led to arrested succession, where soil seed bank became dominated by weedy shrubs over time. Frequent or short jhum cycles impede the ecosystem's ability to recover by subjecting it to repeated 'shocks' that disrupt its physical and biological processes. This Favors only those species capable of thriving in the face of such 'shocks', to the detriment of other flora and fauna.

Observation was also made by Bruun et al.2009 [18] where they found higher growth of pioneer species in fallows due to their shallow root systems and the ability to take up nutrients from the top layer. Studies found that shifting cultivation fallows demonstrate faster recovery of vegetation in terms of basal area, regeneration, and species accumulation compared to other human-modified and abandoned lands, such as pastures, agroforestry sites, and plantations [19].

Another key concern is the deforestation linked to shifting cultivation, leading to carbon emissions and biodiversity loss. However, indigenous farmers often carefully protect their endemic species, which hold ecological value for their communities. The slash-and-burn technique is used to clear forest patches and prepare land for sowing. Research conducted by Levasseur et al.,2000 [20] in Arunachal Pradesh found that not all trees are felled during the clearing process. In fact, the preservation of species like Artocarpus heterophyllus, Terminalia myriocarpa, and Morus levigata while clearing vegetation, is guite common in the shifting cultivation system evolved by Mayan communities in the New World. For weeks, the felled tree branches and other vegetation are left to dry in the open. Just before the spring showers arrive in March, these materials are then purposefully set ablaze, a practice that is often criticized by scientists and experts as contributing to carbon emissions. However, villagers find fire to be the simplest way to remove unwanted vegetation from their farm plots and use the ashes as natural fertilizer to enrich the soil.

Deleterious effects of shifting cultivation such as forest and biodiversity loss, soil erosion, and environmental impacts similar are often consequences of shortened fallow cycles which are likely an offshoot of increasing human population. Therefore, rigorous and sustained investigations are paramount for enhancing our understanding of this issue, which is essential for the restoration and management of tropical forests rebounding from shifting cultivation and other human-induced disruptions. Extending the jhum cultivation cycle to 20-30 years shows promise for mitigating some negative impacts. However, it is crucial to also explore methods that can rapidly restore jhum fallows to forests and support natural soil rejuvenation. Modifying land use practices to enhance productivity is essential for sustainable agriculture. In this regard, traditional agroforestry solutions such as Agroforestry present a highly Homestead suitable approach which is variedly known as Bari in Assamese, Hembiri in Karbi, Nolaihaphai in Dimasa, Intathingri in Singpho, Shong in Jaintia/Pnar and so on. This strategy can strike a balance between agricultural productivity and environmental conservation, providing а sustainable pathway forward for communities engaged in shifting cultivation practices.

2.2 Underappreciation of the Significance of Traditional Customs and Practices Leads to Alienation

A key obstacle facing shifting cultivation is insecurity regarding land tenure. Often,

indigenous and tribal populations lack clearly defined property rights over the lands they farm. This dearth of tenure security can precipitate conflicts. displacements. and impede investments in sustainable land management approaches [21]. The land ownership and tenure system of different tribes of the north-east is highly complicated and diverse. It ranges from private ownership to community ownership [22]. Among the Angamis and Chakhesangs of Nagaland, land ownership and inheritance rights are held individually. In the Naga community, land ownership is divided into distinct categories: family land held by members of a kinship group, privately owned individual plots, and communal clan and village lands managed by the village council of elders or the chief. In contrast, the Ang system among the Semas and Lothas vests land assets in the chieftain. Matrilineal societies like the Khasis, Jaintias, and Garos of Meghalava pass ownership and inheritance to the voungest daughter of the Syiem, Nokma, or Doloi, but management and transfer decisions are made by their related advisers [22].

In Arunachal Pradesh, with no land-use policy and no clear identity of land ownership [23], suggesting any alternatives would just remain academic. The ability to acquire land depends on one's social standing. As a result, only a small number of individuals with extensive land holdings can adopt alternatives to shifting cultivation while maintaining their traditional practices. However, a person with limited land may need to take a risk in this situation, and ultimately, the alternatives may not be well-suited [12]. In Meghalaya, the prevailing land tenure system results in only 1145.19 sq km (5% of total forest areas) of forest areas being directly under the control of the State Forest Department, which includes Reserved Forests. Protected Forests. National Parks, Wildlife Sanctuaries, and Parks and Gardens. The remainder of the forest areas belong to communities, clans, private individuals, and District Councils [24]. And yet, the vital importance and role of these local communities in forest management and governance are consistently undervalued. To transition from shifting cultivation, it is crucial that the shifting cultivators, or Jhumias, are provided with land they can cultivate and generate profits from on a long-term basis [25].

Consequently, the forest department has failed to establish genuine partnerships with local communities, hindering effective forest stewardship. On the other hand, increasing development in the northeastern region has weakened community control and led to privatization, the resulting transformation in land use patterns has compelled villagers facing financial pressures, such as for children's education and medical needs, to relinquish their land in exchange for cash. Various reports indicate that central planning has seldom recognized and respected the significance of traditional customs and practices that are deeply rooted in local cultures and often sustainable. These longstanding traditional practices have provided livelihoods for indigenous populations, yet policymakers have frequently overlooked their value and importance. Therefore, investing agencies, including the government, should carefully consider the long-term social and environmental implications before introducing exotic cash crops.

2.3 Biodiversity and Ecosystem Services in Shifting Cultivation Landscapes

Research by Borah et al., 2022 [26] has uncovered the remarkable biodiversity present within shifting cultivation systems in Northeast India, a well-known biodiversity hotspot. Their findings indicate that active farmland and regenerating fallows support greater species richness than even old-growth forests in the region. While mature forests harbor a unique bird community, including vulnerable and endangered species, the avian communities of regenerating fallows increasingly resemble those of the undisturbed forests as succession progresses. Farmland and regenerating fallows had bird species richness similar to old-growth forests, research on consistent with multi-strata agroforestry systems in India [27] and elsewhere [28]. According to Norgrove et al.,2016 [29] management decisions by local communities can influence the rate of fallow recovery in shifting cultivation, that in turn can lead to varying degrees of biodiversity recovery across sites. Shifting cultivation landscapes in Northeast India hold significant potential for co-benefits. This can be realized by safeguarding old-growth forests from conversion to farmland, permanently abandoning mature secondary forests currently under shifting cultivation, and boosting carbon storage through natural forest regeneration within shiftina cultivation svstem [28.30]. the Shifting cultivation can be a sustainable practice that supports biodiversity when large forest tracts are preserved, few trees are retained during farming, and fallow periods are lengthy [31].

threat is posed by large-scale Another plantations of commercial and perennial crops including rubber and oil palm in lands where shifting cultivation was prevalent [32,33]. The area under plantation crops grew significantly, and betel nut and citrus cultivation doubled and rubber plantations, which were previously absent, had spread across more than 70 km² by 2015 [34,35]. These rapid, large-scale changes in land use have strongly contributed to declines in agrobiodiversity and the ecosystem services that support the region, with both long-term and short-term consequences for food security [4]. The conversion of fallow forest lands into commercial plantations by tribal communities poses a concerning risk of inadvertently promoting monoculture, which could come at the expense of biodiversity and ecological wellbeing. This shift in land use, if left unaddressed, could have detrimental consequences for the delicate balance of the region's diverse and fragile ecosystems. Yet shifting cultivation is regarded as one of the major drivers of deforestation and forest degradation in this region and hence is discouraged in local landuse policies [36,14].

Numerous wild plants, with medicinal and nutritional value as well as cultivated crop varieties, thriving in shifting production system. A study conducted by Pandev et al., 2022 [37] in the Garo Hills region found that the food system biodiversity within the jhum landscape is strongly associated with dietary diversity among the Garo community. Nongrum et al. 2021 [38] observed that Garo women have higher levels of dietary diversity. They play a pivotal role in preserving the rich biodiversity by conserving the seeds of traditional crops [27]. However, the extensive changes in land use within the ihum landscape of Garo Hills, Meghalaya, have shifted the primary focus of farming to cash crops [34,35]. This monoculture-based approach has transformed food consumption patterns, ushering in a new food system that leaves both farmers and consumers vulnerable to market fluctuations [4].

Therefore, bolstering local food systems is a crucial priority to minimize external dependencies and vulnerabilities. This can be achieved by championing traditional, nutrient-dense crop varieties that are also resilient to climate-related stresses. The prioritization of monoculture crops for short-term financial gains poses a grave threat to the food security, nutritional well-being, and livelihood stability of low-income households who rely on the resilient and climate-adaptive

practices of jhum cultivation. The introduction of hybrid and genetically modified high-yielding commercial crops has led to the neglect of indigenous crop varieties, with many traditional cultivars disappearing from the landscape. These native crops possess inherent advantages, such as resilience to climate change and resistance to pests, in contrast to the high-yielding varieties that require chemical inputs and cannot withstand environmental stresses. From a socioeconomic standpoint, the preservation and conservation of these traditional crop varieties is paramount, as they enable self-sufficient farming without dependence on external inputs. Recognizing the importance of indigenous agriculture, the Government of India has implemented incentives and awards, such as the Plant Genome Saviour Award, to encourage local farmers to cultivate and conserve their native crop diversity. Consequently, policies that support and enhance the viability of jhum cultivation can address the food security needs of indigenous communities while preserving their traditional ecological knowledge and practices.

2.4 The Legacy and Struggles of Shifting Cultivation in India's Agricultural Policies

Shifting cultivation, which is more akin to agricultural practices, remains under the purview of the Ministry of Forest and Environment due to its colonial-era legacy, with limited involvement the Ministry of Agriculture. from India's agricultural policies and initiatives have largely overlooked and failed to support shifting cultivation practices. According to Darlong 2019 [39] these traditional farming methods are often sidelined and not regarded as integral components of mainstream agriculture. This has led to the perception that shifting cultivation takes place on forested lands. British colonial forest policies in India sought to assert state dominance the interests of local communities, over safeguarded particularly those who and managed forests based on their traditional practices and shifting cultivators were consistently targeted in this process [39]. Colonial rulers regarded it as a detrimental landpractice. associating it with forest use degradation and diminished revenue. This outdated mindset has persisted, shaping policies that continue to criticize shifting cultivation. Such perceptions have led to the policies that altogether ruled out traditional agriculture through legal enforcements (e.g., establishment of national parks) or provided economic incentives

to people for adoption of modern agrotechnologies [40].

A five-country study initiated by International Center for Integrated Mountain Development (ICIMOD). А mountain research and development think tank based in Kathmandu, Nepal, observed that 'indigenous farmers have much to teach the world about the efficient use of their landscape for combined agriculture and Notably, the traditional practices forestrv'. maintain robust crop yields, prevent soil degradation, and foster biodiversity conservation - lessons that the world would be wise to heed [5]. This study made the government of India to pussyfoot and come up with a new policy document, released by NITI Aayog in 2018 titled, "Shifting Cultivation: Towards a Transformational Approach." However, according to authors of NITI Aayog Report on Jhum, Special Issue, 2023, five years have passed by since the document was released and there seems to be no action.

Jhum cultivation persists largely due to the absence of feasible alternative technologies. Several reports indicate that the farming system models developed by the Indian Council of Agricultural Research (ICAR) regional center were ill-suited to the region's diverse terrain and altitudinal conditions. A key factor is that the hillocks are generally not owned by a single individual, but rather shared among multiple families. The reports also assess alternative land use interventions suggested by the Indian Council of Agricultural Research to replace cultivation [22]. However, simply shifting replacing jhum cultivation is unlikely to succeed, and a more prudent approach would be to build upon and enhance this existing practice. However, externally supported initiatives in Northeast India, such as those financed by the India-Canada Environment Facility and the International Fund for Agricultural Development, seek to enhance shifting cultivation areas and livelihoods. The North Eastern Region Community Resource Management Project for Upland Areas (NERCORMP) project focus on adapting traditional jhum cultivation, encouraging more profitable crops, and converting jhum lands into community-conserved areas for sustainable harvesting of non-timber forest products. Another initiative is the Nagaland Environmental Protection and Economic Development Project (NEPED) project in Nagaland, which was the launched under Indo-Canadian Environmental Facility Program. Initially targeting around 1200 villages, the project has now streamlined its efforts to a more manageable scale. The aim is to link these activities with marketing opportunities and foster small-scale entrepreneurship leveraging the local natural resource base and ecological conditions. This process has embarked on addressing the economic and environmental challenges associated with jhum cultivation.

In the current market-driven landscape, these traditional practices require complementary modern inputs to fortify and maximize their productivity. Such interventions can be highly effective in mitigating regional economic disparities and improving the livelihoods of marginalized upland farmers. It is imperative to leverage traditional crop resources and deploy appropriate scientific solutions to drive advancements in agriculture. Not necessarily all the traditional ecological knowledge may be good, and therefore, we need to validate the traditional ecological knowledge through which scientific empirical analysis, the community is accustomed to, and then integrate the validated knowledge into formal academic and policy frameworks. This approach can help ensure that traditional practices are complemented with modern scientific inputs to enhance their effectiveness and contribute to more sustainable and equitable agricultural development.

3. CONCLUSION AND RECOMMENDA-TIONS

Despite the global changes intruding into the socio-economic sphere of the tribal communities across Northeastern region of India, shifting cultivation continues to play an important role in providing livelihoods and food security to the rural tribal households. It seems that shifting cultivation is closely tied to the cultural identity of the tribal people. Therefore, its importance lies beyond mere economic concerns. Though with government interventions and under innovative shifting cultivation, the farmers in the region have switched to newer methods of cultivation, shifting cultivation accommodating at the same time the value system and needs of the tribal society.

The contribution of jhum to loss of forest capital may not be as severe as that of other factors such as logging or conversion of natural forests to plantations [40]. The study by Geist et al. in 2002 [41] suggests that complex interactions between multiple factors, rather than simplistic single-factor explanations, drive deforestation.

The analysis indicates that public and individual choices at the fundamental level are largely influenced by shifting economic opportunities and policies at national and global scales, mediated by local institutional factors, rather than just being attributed to shifting cultivators and population growth. At the proximate level, regionally distinct agricultural expansion, wood extraction. dam building and massive infrastructure development contribute to deforestation whose devastating impacts are often ignored.

While the negative impacts of shifting cultivation cannot be ignored, it is important to consider the opposing perspective. Following a shifting cultivation cycle, the secondary growth forests can recover up to 80% of their tree species diversity [42] and more than 50% of their aboveground biomass [43] within a relatively short timeframe of approximately 20-30 years. This demonstrates the resilience of these ecosystems and their ability to regenerate. Furthermore, the practice of shifting cultivation is deeply rooted in the local traditions and sociocultural values of the indigenous populations, providina essential livelihoods. Mounting evidence from research across the tropical and sub-tropical forest regions of Asia, central Africa, and South America - where shifting agriculture is prevalent practice - demonstrates that а indigenous farming systems effectively maintain high levels of biodiversity. These systems not only ensure food and nutritional security but also provide vital livelihoods for populations living in tropical forest areas.

Land-use conversions of forests for timber logging, monoculture cash crops, and infrastructure projects like hydroelectric dams typically benefit stakeholders who do not reside in the local landscape, unlike subsistence-based cultivation [44]. Alternatives shifting often proposed, such as intensive sedentary farming, monocultures, or agroforestry, may potentially have long-term harmful impacts on both farmers and forests.

Considering the intricate nuances and intricacies of the diverse local cultures and cultural practices, any interventions or initiatives must be carefully designed to align with the traditional frameworks to avoid alienation of the local communities and instead enhance their acceptance and ownership of such efforts. The traditional practices are deeply rooted in the socio-cultural fabric of the region and carry significant meaning and significance for the indigenous populations. Therefore, maintaining fidelity to these time-honored traditions is crucial to ensure the long-term sustainability and effectiveness of any proposed changes or alternative approaches. By viewing land use practices dynamic, interconnected as components of cultural and village landscapes, rather than as isolated and unchanging systems, we can gain deeper insights and address knowledge gaps. Introducing formal knowledge into community systems is a valuable endeavor, but it is crucial to first assess the community's receptiveness. For instance, if the community believes in burning practices for soil fertility, it is essential to educate them on alternative, more sustainable methods of maintaining soil fertility without the need for burning. The most effective approach is to facilitate a thought process within the community, rather than simply imposing knowledge from the outside.

The proposed approach should be to extend the jhum cultivation cycle to 20-30 years, while simultaneously reducing the overall jhum area through more sustainable land use choices. These alternative solutions should build upon traditional practices and indigenous knowledge. such as homestead agroforestry, rather than promoting market-driven monocultures like oil palm or exotic timber plantations, which can trap farmers in unfavorable situations. The goal should not be to eliminate jhum cultivation, but to enhance it as a viable and environmentally conscious livelihood option for the local communities. The traditional mountain farmers prioritize local production-driven food selfsufficiency, low-input organic traditional agrofood ecosystems for security. and environmentally friendly approaches to land-use practices like improved management of jhum fallows and diversification of economic avenues to cope with environmental uncertainties and mitigate risks. Therefore, there is an urgent need to redevelop the largely fragmented landscapes of the region in a way that addresses their concerns for sustainable livelihoods, as neglecting these could have serious implications for human security in the region. Promoting the cultivation of indigenous, climate-resistant crop varieties is crucial for fortifying local food systems and minimizing dependence on external inputs. Policymakers must prioritize these validate sustainable practices, traditional knowledge through empirical research, and invest in complementary modern techniques to ensure the long-term agricultural viability and food security of indigenous communities.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

- 1. Borthakur DN. Agriculture of the North Eastern Region. Guwahati: BeeCee Prakashan; 1992.
- 2. Bos SPM, Cornioley T, Dray A, Waeber PO, Garcia CA. Exploring livelihood strategies of shifting cultivation farmers in Assam through games. Sustainability. 2020;12(6):2438.

DOI: 10.3390/su12062438.

- Rahman SA, Rahman MF, Sunderland TCH. Causes and consequences of shifting cultivation and its alternatives in the hill tracts of eastern Bangladesh. Agrofor Syst. 2012;84(2):141-55. DOI: 10.1007/s10457-012-9494-3.
- Behera RN, Nayak DK, Andersen P, Måren IE. From jhum to broom: Agricultural land-use change and food security implications on the Meghalaya Plateau, India. Ambio. 2016;45(1):63-77. DOI: 10.1007/s13280-015-0691-3.
- Kerkhoff E, Sharma E. Debating shifting cultivation in the Eastern Himalayas: Farmers' innovations as lessons for policy.
 G.P.O. Box 3226, Kathmandu, Nepal. International Centre for Integrated Mountain Development (ICIMOD); 2006.
- Raj S. Traditional knowledge, innovation systems and democracy for sustainable agriculture: A case study on Adi tribes of Eastern Himalayas of North-East India. In: Innovation and Sustainable Development in Agriculture and Food. 2010;1-10.
- Manikant Kumar. Environmental impacts and sustainability of shifting cultivation practices in India. International Journal of Research Publication and Reviews. 2023;4(9):1030-6.
- Kalita H, Baruah MS, Datta D, Jini D, Alone RA. Status of shifting cultivation (jhum) in Arunachal Pradesh, India. In: Prakash N, et al., editors. Jhum

improvement for sustaining farm, livelihood and natural resource conservation in the North Eastern Hill Region: Vistas and frontiers. India: ICAR Research Complex for NEH Region; 2017.

- 9. Das D. Demystifying the myth of shifting cultivation: Agronomy in the North-East. Economic and Political Weekly. 2006;41(47):4912-7.
- 10. Hazarika S. Options for natural resource conservation for jhum improvement in NEH Region. In: Prakash N, et al., editors. Jhum improvement for sustaining farm, livelihood and natural resource conservation in the North Eastern Hill Region: Vistas and frontiers. India: ICAR Research Complex for NEH Region; 2017.
- 11. Bawa KS, Dayanandan S. Climate change and biodiversity: Implications for conservation in the tropics. Climatic Change. 1997;39:473-85.
- Arunachalam A, Khan ML, Arunachalam K. Balancing traditional jhum cultivation with modern agroforestry in Eastern Himalaya – A biodiversity hotspot. Current Science. 2002;83(2):117-8.
- Murty D, Kirschbaum MUF, McMurtrie RE, McGilvray H. Does conversion of forest to agricultural land change soil carbon and nitrogen? A review of the literature. Global Change Biology. 2002;8:105-23.
 - DOI: 10.1046/j.1354-1013.2001.00459.x.
- Mukul SA, Herbohn J, Ferraren A, Congdon R. Limited role of shifting cultivation in soil carbon and nutrients recovery in regenerating tropical secondary forests. Frontiers in Environmental Science. 2022;10. DOI: 10.3389/fenvs.2022.1076506.
- Mukul SA, Herbohn J, Firn J, Gregorio N. Carbon and biodiversity outcomes under divergent management scenarios – lessons from upland Philippines shifting cultivation landscapes. In: Cairns M, editor. Farmer innovations and best practices by shifting cultivators in Asia-Pacific. 2021;408-20.
- Funakawa S, Makhrawie M, Pulunggono HB. Soil fertility status under shifting cultivation in East Kalimantan with special reference to mineralization patterns of labile organic matter. Plant and Soil. 2009;319(1):57-66.
- Raharimalala O, Buttler A, Ramohavelo CD, Razanaka S, Sorge JP, Gobat JM. Soil-vegetation patterns in secondary slash and burn successions in Central Menabe,

Madagascar. Agriculture, Ecosystems & Environment. 2010;139(1-2):150-8. DOI: 10.1016/j.agee.2010.07.013.

 Bruun TB, de Neergaard A, Lawrence D, Ziegler AD. Environmental consequences of the demise in swidden cultivation in Southeast Asia: Carbon storage and soil quality. Human Ecology. 2009;37(3):375-88.

DOI: 10.1007/s10745-009-9257-y.

- Ferguson BG, Vandermeer J, Morales H, Griffith DM. Post-agricultural succession in El Petén, Guatemala. Conservation Biology. 2001;17(3):818-28.
- Levasseur V, Olivier A. The farming system and traditional agroforestry systems in the Maya community of San Jose, Belize. Agroforestry Systems. 2000;49(3):275-88. DOI: 10.1023/A:1006327403980.
- 21. Fernandes W. Shifting cultivation and environmental change: Indigenous people, agriculture, and forest conservation. Oxford: Oxford University Press; 2008.
- Choudhury D, Sundriyal RC. Factors contributing to the marginalization of shifting cultivation in North-East India: Micro-scale issues. Outlook on Agric. 2003;32(1):17-28. DOI: 10.5367/00000003101294226.
 - DUI: 10.5367/00000003101294226.
- Bakshi PM, Kusum. Land system of Arunachal Pradesh. Mumbai: N. M. Tripathi Pvt Ltd; 1989.
- Forest Survey of India. State of Forest Report 2017. Available:https://www.megforest.gov.in/for est_cover.html.
- 25. Darlong VT. To jhum or not to jhum: Policy perspective on shifting cultivation. The Missing Link (TML). Society for Environment and Communication, India. 2004;143.
- 26. Borah JR, Gilroy JJ, Evans KL, Edwards DP. The value of shifting cultivation for biodiversity in Northeast India. Divers Distrib. 2022. doi: 10.1111/ddi.13605.
- Ellena R, Nongkynrih KA. Changing gender roles and relations in food provisioning among matrilineal Khasi and patrilineal Chakhesang Indigenous rural people of North-East India. Matern Maternal & Child Nutrition. 2017;13: e12560.
- 28. Basham EW, González del Pliego P, Acosta-Galvis AR, Woodcock P, Medina Uribe CA, Haugaasen T, et al. Quantifying carbon and amphibian co-benefits from

secondary forest regeneration in the Tropical Andes. Animal Conservation. 2016;19(6):548-60.

DOI: 10.1111/acv.12276.

- 29. Norgrove L, Beck J. Biodiversity function and resilience in tropical agroforestry systems, including shifting cultivation. Current Forestry Reports. 2016;2(1):62-80.
- Gilroy JJ, Woodcock P, Edwards FA, Wheeler C, Baptiste BLG, Medina Uribe CA, et al. Cheap carbon and biodiversity co-benefits from forest regeneration in a hotspot of endemism. Nature Climate Change. 2014;4(6):503-7. DOI: 10.1038/nclimate2200.
- N'Dja K, Justin K, Guillaume D. Successional patterns of plant species and community diversity in a semi-deciduous tropical forest under shifting cultivation. Nature Climate Change. 2008;19(6):809-20.
- Ziegler AD, Phelps J, Yuen JQ, Webb EL, Lawrence D, Fox JM, et al. Carbon outcomes of major land-cover transitions in Southeast Asia: Great uncertainties and REDD+ policy implications. Global Change Biology. 2012;18(9):3087-99.
 DOI: 10.1111/j.1365-2486.2012.02747 x

DOI: 10.1111/j.1365-2486.2012.02747.x.

- Terefe B, Kim DG. Shifting cultivation maintains but its conversion to monocropping decreases soil carbon and nitrogen stocks compared to natural forest in Western Ethiopia. Plant and Soil. 2020;453(1-2):105-17. DOI: 10.1007/s11104-019-03942-0.
- 34. Government of Meghalaya. Statistical Handbook. Shillong: Directorate of Economics and Statistics; 2019.
- National Remote Sensing Centre. Wasteland Atlas of India. Department of Space; 2019. Available:https://dolr.gov.in/documents/wa steland-atlas-of-india.
- Hett C, Castella J, Heinimann A, Messerli P, Pfund J. A landscape mosaics approach for characterizing swidden systems from a

REDD+ perspective. Applied Geography. 2012;32:608-18.

DOI: 10.1016/j.apgeog.2011.07.011.

37. Pandey DK, Momin KC, Dubey SK, et al. Biodiversity in agricultural and food systems of jhum landscape in the West Garo Hills, North-eastern India. Food Sec. 2022;14:791-804.

DOI: 10.1007/s12571-021-01251-y.

- Nongrum MS, Pawera L, Mawroh B. Dietary diversity and its determinants among Khasi and Garo indigenous women (15 to 49 years) in Meghalaya, northeast India. Nutrition and Health. 2021;02601060211016629. DOI: 10.1177/02601060211016629.
- Darlong V. Reflections on the impacts of state policies on shifting cultivators in Northeast India. In: Shifting cultivation policies: Balancing environmental and social sustainability. 2019. p. 344-78. DOI: 10.1079/9781786391797.0344.
- 40. Shimrah T, Rao KS, Saxena KG. The shifting agricultural system (jhum) and strategies for sustainable agroecosystems in Northeast India. Agroecology and Sustainable Food Systems. 2015;39(2): 155-77.

DOI: 10.1080/21683565.2015.1011173.

- 41. Geist HJ, Lambin EF. Proximate causes and underlying driving forces of tropical deforestation. BioScience. 2002;52(2):143-50.DOI:10.1641/0006-3568(2002)052[0143:PCAUDF]2.0.CO;2.
- 42. Rozendaal DMA, Bongers F, Aide TM, et al. Biodiversity recovery of Neotropical secondary forests. Science Advances. 2019;5:eau3114.

DOI: 10.1126/sciadv.aau3114.

- Poorter L, Bongers F, Aide TM, et al. Biomass resilience of Neotropical secondary forests. Nature. 2016;530:211-4. DOI: 10.1038/nature16512.
- 44. Dressier W. Disentangling Tagbanua lifeways, swidden and conservation on Palawan Island. Human Ecology. 2005;12(1):21-9.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/128483